



H. Lundbeck A/S

Ottiliavej 9 DK-2500 Valby, Danmark Tlf. +45 36 30 13 11 Fax +45 36 43 82 62 E-mail information@lundbeck.com www.lundbeck.com

Press Release

Valby, December 10, 2014

Otsuka and Lundbeck present Phase III data on brexpiprazole as adjunctive therapy in adult patients with major depression at the American College of Neuropsychopharmacology annual meeting

- Results from two Phase III clinical studies demonstrated the effects of brexpiprazole
 as adjunctive treatment in patients with major depressive disorder (MDD) who had an
 inadequate response to monotherapy antidepressant treatments (ADT).
- Brexpiprazole is a serotonin-dopamine activity modulator (SDAM) and is believed to
 possess a balanced combination of potent activities at multiple receptors in the brain
 including partial agonist activity at dopamine D2 and serotonin 5HT1A receptors, and
 antagonist activity at serotonin 5HT2A receptors and noradrenergic alpha1B/2C
 receptors.
- Also being presented at ACNP are results from two phase III clinical studies of brexpiprazole in adult patients with schizophrenia.

Otsuka Pharmaceutical Co., Ltd. (Otsuka) and H. Lundbeck A/S (Lundbeck) today announced the presentation of Phase III study results evaluating the efficacy of investigational compound brexpiprazole as adjunctive treatment to antidepressant therapy (ADT) in patients with major depressive disorder (MDD) at the 53rd Annual Meeting of the American College of Neuropsychopharmacology (ACNP) in Phoenix, Arizona. The data were shared in a poster presentation, "Efficacy and Safety of Adjunctive Brexpiprazole (OPC-34712) in Major Depressive Disorder: Results of Two Pivotal Clinical Studies."

"Inadequate response to monotherapy antidepressant treatment is frustrating for patients and the likelihood of remission with multiple lines of therapy decreases substantially with each successive treatment," said Michael E. Thase, MD, Professor of Psychiatry, Director, Mood and Anxiety Program, University of Pennsylvania School of Medicine and study investigator. "These results are very encouraging in that they provide evidence of efficacy and safety of brexpiprazole as an adjunctive treatment to antidepressant therapy in patients with major depressive disorder who had an inadequate response to antidepressant therapy."

MDD Study Results

The poster featured results of two Phase III clinical studies evaluating the efficacy, safety and tolerability of adjunctive brexpiprazole in patients with MDD and inadequate response to





ADT (Study 1: NCT01360645; Study 2: NCT01360632). Patients with MDD who failed to reach adequate response during 1-3 treatment attempts with ADT (which is commonly found in current clinical practice) were enrolled and received an additional trial with a (single-blind) ADT for 8 weeks. Those patients who still failed to reach an adequate response throughout this phase were then randomized (double-blind) to ADT and brexpiprazole or ADT and placebo for 6 weeks. The primary endpoint for both studies was change in MADRS (Montgomery–Åsberg Depression Rating Scale) Total Score from baseline to Week 6. Prespecified analyses were conducted both on the efficacy population and on the final protocol population (fulfilling amended randomization criteria) for each individual study.

In both studies, key findings included:

- Adjunctive brexpiprazole showed greater improvement than adjunctive placebo in MADRS total score at Week 6 in the efficacy population per final protocol in Study 1 (2mg+ADT [N=175]: -3.21, p=0.0002), and in Study 2 (1mg+ADT [N=211]: -1.30, p=0.0737; 3mg+ADT [N=213]: -1.95, p=0.0079). Similar results were observed for the efficacy population in both studies.
- The completion rate was high (>90%) and comparable across brexpiprazole and placebo groups. Discontinuations due to adverse events were low across all groups (1mg = 1.3%, 2mg = 3.2%, 3mg = 3.5%, placebo = 0.7%) and only one patient discontinued due to lack of efficacy (in the brexpiprazole 1mg group).
- All doses of adjunctive brexpiprazole resulted in notably low levels of sedating or activating side effects.
- The most frequent adverse events (incidence >5% in any group and more than twice the incidence in the placebo group across the two studies) included akathisia (4.4%, 7.4%, 13.5% vs.1.7%), weight increase (6.6%, 8.0%, 5.7% vs. 1.9%), tremor (4.0%, 2.1%, 5.2% vs. 2.2%) somnolence (4.0%, 4.3%, 5.7% vs. 0.5%) and nasopharyngitis (6.6%, 1.1%. 3.1% vs. 1.7%), in the brexpiprazole 1mg+ADT (N=226), 2mg+ADT (N=188), 3mg+ADT (N=229) versus combined placebo + ADT groups (N=411), respectively.

The results from Study 1 were previously presented in a poster session at the 22nd European Psychiatry Association Congress (EPA) in March 2014.

"These results support previously reported data investigating the effect of brexpiprazole for patients with major depressive disorder," said William Carson, MD, CEO, Otsuka Pharmaceutical Development & Commercialization, Inc. "There are more than 14 million adults with this condition in the U.S., of which a significant portion continuously suffer from inadequate response to antidepressant therapy, and new therapeutic options are needed to help those patients struggling to find effective, tolerable treatments. These study results suggest we are on the right track."

"We are committed to patients who suffer from depression," said Anders Gersel Pedersen, MD, EVP and head of R&D in Lundbeck. "We are very grateful to patients, investigators and the broader mental health community for their collaboration and support of our extensive





clinical program studying brexpiprazole, and we believe in its potential to make a difference for patients and their families affected by major depressive disorder."

Otsuka and Lundbeck will also present at ACNP Phase III data results evaluating the effect of brexpiprazole as a monotherapy in adult patients with schizophrenia. The data will be shared in two poster presentations, "A Multicenter, Randomized, Controlled, Phase III Trial of Fixed-dose Brexpiprazole for the Treatment of Adults with Acute Schizophrenia" and "Brexpiprazole for the Treatment of Acute Schizophrenia: A Randomized, Controlled Trial."

About Brexpiprazole (OPC-34712)

Brexpiprazole is a novel investigational psychotropic compound discovered by Otsuka and under co-development with Lundbeck. Brexpiprazole is a serotonin-dopamine activity modulator (SDAM) that acts as a partial agonist at 5-HT1A and dopamine D2 receptors, and an antagonist at 5-HT2A and noradrenaline alpha1B/2C receptors, all with similar high potency (< 1nM). A New Drug Application for brexpiprazole has been filed with the US FDA and the PDUFA date is in July 2015.

Lundbeck Contacts

Mads Kronborg, Director, Media Relations Telephone (direct): +45 36 43 28 51 Lars Otto Andersen-Lange Telephone (direct): +45 30 83 26 57

U.S. Ashleigh Duchene aduc@lundbeck.com +1 847 282 1164

Otsuka Contacts

Investors: Yoko Ishii Otsuka Holdings Co., Ltd. Ishiiyo@Otsuka.jp +81 3 6361 7411

Media:
JAPAN/ASIA
Jeffrey Gilbert
Leader, Pharmaceutical PR
Otsuka Pharmaceutical Co, Ltd.
gilbert.jeffrey@otsuka.co.jp
+81 3 6361 7379, +81 80 8728 6039

U.S.
Rose Weldon
Corporate Communications
Otsuka America Pharmaceuticals
rose.weldon@otsuka-us.com
+1 609 524 6879

About Lundbeck

H. Lundbeck A/S (LUN.CO, LUN DC, HLUYY) is a global pharmaceutical company specialized in brain diseases. For more than 50 years, we have been at the forefront of research within neuroscience. Our key areas of focus are alcohol dependence, Alzheimer's





disease, bipolar disorder, depression/anxiety, epilepsy, Huntington's disease, Parkinson's disease, schizophrenia, stroke and symptomatic neurogenic orthostatic hypotension (NOH).

An estimated 700 million people worldwide are living with brain disease and far too many suffer due to inadequate treatment, discrimination, a reduced number of working days, early retirement and other unnecessary consequences. Every day, we strive for improved treatment and a better life for people living with brain disease – we call this Progress in Mind.

Read more at www.lundbeck.com/global/about-us/progress-in-mind.

Our approximately 6,000 employees in 57 countries are engaged in the entire value chain throughout research, development, production, marketing and sales. Our pipeline consists of several late-stage development programmes and our products are available in more 100 countries. We have research centres in China, Denmark and the United States and production facilities in China, Denmark, France and Italy. Lundbeck generated revenue of approximately DKK15.3 billion in 2013 (EUR2.1 billion; USD2.7 billion).

For additional information, we encourage you to visit our corporate site www.lundbeck.com.

About Otsuka Pharmaceutical Co., Ltd.

Otsuka Pharmaceutical is a global healthcare company with the corporate philosophy: "Otsuka – people creating new products for better health worldwide." Otsuka researches, develops, manufactures and markets innovative and original products, with a focus on pharmaceutical products for the treatment of diseases and nutraceutical products for the maintenance of everyday health.

In pharmaceuticals, Otsuka is a leader in the challenging area of mental health and also has research programs on several under-addressed diseases including tuberculosis, a significant global public health issue. These commitments illustrate how Otsuka is a "big venture" company at heart, applying a youthful spirit of creativity in everything it does.

Otsuka Pharmaceutical, which employs approximately 28,700 people worldwide, is a wholly owned subsidiary of Otsuka Holdings Co., Ltd., the holding company for the Otsuka Group that is headquartered in Tokyo, Japan. The Otsuka Group has business operations in 26 countries and regions around the world, with consolidated sales of approximately USD 14.1 billion for fiscal year 2013 (4/1/2013-3/31/2014.) Otsuka welcomes you to visit its global website at https://www.otsuka.co.jp/en.